

Comment on “The double–Kerr solution” by W B Bonnor and B R Steadman

V S Manko[†] and E Ruiz[‡]

[†]Departamento de Física,
Centro de Investigación y de Estudios Avanzados del IPN,
A.P. 14-740, 07000 México D.F., Mexico

[‡]Area de Física Teórica, Universidad de Salamanca,
37008 Salamanca, Spain

Abstract

Erroneous statements on the mathematical and physical properties of the double–Kerr solution made in a recent paper (2004 *Class. Quantum Grav.* **21**, 2723) are commented.

PACS numbers: 0420

The well-known Kramer–Neugebauer (KN) solution [1] for two Kerr black holes [2] has been analyzed in a recent work [3] by means of an approximation procedure. Since the paper [3] contains various erroneous statements on the properties of the KN spacetime, some comments on it seem appropriate.

(i) First of all, it is worthwhile pointing out that the BS approximate potential (32) of [3] lacks generality, that invalidates the physical conclusions which Bonnor and Steadman believe to have been drawn for the whole KN spacetime. This is because their statement following formula (32) – “*As expected, the solution has five parameters: m_s , q_s , b* ” – is wrong. The general asymptotically flat subfamily of double–Kerr spacetimes contains *six* arbitrary real parameters, and the absence of one parameter in the BS scheme is

explained by the restrictive character of the reparametrization (12)–(15) employed in [3] instead of the original KN parametrization [1]. Indeed, Kramer and Neugebauer used in their paper *eight* real constants K_l , ω_l , $l = 1, 2, 3, 4$, of which the constants K_l can be subjected to the constraint $K_1 + K_2 + K_3 + K_4 = \text{const}$ due to the liberty in the shift along the z -axis, while the four constants ω_l give rise to the unphysical NUT parameter which can be eliminated by an appropriate unitary transformation. Therefore, in the general asymptotically flat case one is left with *six* arbitrary real constants (8 parameters minus 2 constraints), and not *five* as in [3]. A correct reparametrization of the KN solution which does not lead to the loss of one parameter was performed in [4].

Below we give a possible 6-parameter axis expression for the Ernst potential [5] of the asymptotically flat double-Kerr solution:

$$\mathcal{E}(\rho = 0, z) = \frac{z - b - M_1 - i(A_1 + \nu)}{z - b + M_1 - i(A_1 - \nu)} \cdot \frac{z + b - M_2 - i(A_2 - \nu)}{z + b + M_2 - i(A_2 + \nu)}, \quad (1)$$

where the parameters M_i represent the masses of the Kerr particles, A_i their angular momenta per unit mass, b is the separation constant, and ν can be associated with the angular momentum of the part of the intermediate region.

(ii) It is important to clarify that Bonnor and Steadman expand the KN solution not only in powers of masses, as they claim, but also in powers of the angular momenta per unit mass since the latter are implicitly defined in terms of the mass parameters. This fact is a key point for understanding why the approximate potential they obtained describes a specific system of black holes, and not a system involving superextreme Kerr constituents.

(iii) It seems that the biggest confusion of the paper [3] is connected with the physical interpretation of the region separating the particles in the double-Kerr solution. Bonnor and Steadman take as granted that the black holes of the KN solution are separated by a massless strut (see Fig. 1a), most probably being unaware of the fact that the parameters they are using have a rather formal character (their “masses m_1, m_2 ” for instance are not really the masses of black holes) and also describe the systems of not separated Kerr constituents, e.g., as shown in Fig. 1b where a “massive strut” is present (the two rods can also intersect partially). Moreover, one does not know from the very beginning which situation describes the KN solution, so that the correct choice of values of the parameters is of paramount importance.




Figure 1: Possible configurations of two black holes in the KN solution.

Analyzing the case of two separated black holes, one should remember that after having warranted the asymptotic flatness of the KN solution, the next priority is to achieve

$$\omega = 0 \quad (2)$$

on the part $K_3 < z < K_2$ of the z -axis, the function ω entering the axisymmetric line element

$$ds^2 = f^{-1}[e^{2\gamma}(d\rho^2 + dz^2) + \rho^2 d\varphi^2] - f(dt - \omega d\varphi)^2 \quad (3)$$

(the other two metric coefficients are f and γ). Fulfillment of the condition (2) converts the region $\rho = 0$, $K_3 < z < K_2$ into a *massless line strut* containing no closed timelike curves (that is why (2) is sometimes called the axis condition [4]).

The case $\omega \neq 0$ for $K_3 < z < K_2$ describes the systems of two overlapping Kerr subextreme constituents of the type shown in Fig. 1b. In this case the intermediate region $\rho = 0$, $K_3 < z < K_2$ possesses in general a non-zero mass and a non-zero angular momentum.

In view of the above said, the physical interpretation given in [3] to the solution I – two Kerr particles supported by a *massless* strut – is erroneous

because $\omega \neq 0$ between the particles and hence the intermediate region in that solution is not massless. Consequently, interpretation of the latter region as “a massless spinning rod with angular momentum $2m_1m_2q$ ” is erroneous too, and this can be easily illustrated by calculating the Komar mass M of any part of that rod with the aid of the formula [6]

$$M = -\frac{1}{4}\omega[\psi(z = z_1) - \psi(z = z_2)], \quad z_1 > z_2, \quad (4)$$

where the functions ψ and ω are defined by the formulae (38), (41) and (45) of [3] taken at $\rho = 0$, $|z| < b$, and z_1, z_2 are two arbitrary points on the rod. Thus, for instance, choosing $m_1 = m_2 = 1$, $b = 3$, $q = 0.6$ in the latter formulae, one obtains from (4) that $M \approx -18.088$ for $z_1 = 2.5$, $z_2 = -2.5$.

Therefore, the BS solution I belongs to the case of two overlapping subextreme Kerr constituents (Fig. 1b) and its intermediate region has some internal structure which gives rise to the closed timelike curves due to the presence of the negative mass. This solution is physically meaningless and is inapplicable for the description of two separated particles. At the same time, it cannot of course cast any doubt on the importance of the general KN solution since one should restrict his consideration to the physical range of the parameters and exclude unphysical configurations.

There was a search for the equilibrium states of two Kerr particles since 1980, and progress was achieved on the way of extending the original KN solution [1] to the case involving superextreme Kerr constituents [4]. At present the double-Kerr balance problem can be considered as completely solved [7, 8], and the paper [3] in no way contributes either into the analysis of the genuine equilibrium configurations of two spinning particles without a strut, or even into shedding some new light on the systems composed of two Kerr black holes.

References

- [1] Kramer D and Neugebauer G 1980 *Phys. Lett. A* **75** 259
- [2] Kerr R P 1963 *Phys. Rev. Lett.* **11** 237
- [3] Bonnor W B and Steadman B R 2004 *Class. Quantum Grav.* **21** 2723
- [4] Dietz W and Hoenselaers C 1985 *Ann. Phys. (NY)* **165** 319

- [5] Ernst F J 1968 *Phys. Rev.* **167** 1175
- [6] Tomimatsu A 1983 *Prog. Theor. Phys.* **70** 385
- [7] Manko V S and Ruiz E 2001 *Class. Quantum Grav.* **18** L11
- [8] Manko V S and Ruiz E 2002 *Class. Quantum Grav.* **19** 3077